

Approximation of Riemann's Zeta Function by Finite Dirichlet Series. II

YU. V. MATIYASEVICH

Steklov Institute of Mathematics at St.Petersburg, Russia

<http://logic.pdmi.ras.ru/~yumat/personaljournal/finitedirichlet>

Zeta zeroes are very knowledgable

Zeta zeroes are very knowledgable

Zeta zeroes "know about"

- ▶ prime numbers (via von Mangoldt's theorem)

Zeta zeroes are very knowledgable

Zeta zeroes "know about"

- ▶ prime numbers (via von Mangoldt's theorem)
- ▶ the initial trivial zeroes

Zeta zeroes are very knowledgable

Zeta zeroes "know about"

- ▶ prime numbers (via von Mangoldt's theorem)
- ▶ the initial trivial zeroes
- ▶ other non-trivial zeroes

Zeta zeroes are very knowledgable

Zeta zeroes "know about"

- ▶ prime numbers (via von Mangoldt's theorem)
- ▶ the initial trivial zeroes
- ▶ other non-trivial zeroes
- ▶ the pole of the zeta function


Zeta zeroes are very knowledgable

Zeta zeroes "know about"

- ▶ prime numbers (via von Mangoldt's theorem)
- ▶ the initial trivial zeroes
- ▶ other non-trivial zeroes
- ▶ the pole of the zeta function via the zeroes of the factor $1 - 2 \cdot 2^{-s}$ cancelling the pole

Theorem of von Mangoldt

$$\psi(x) = x - \sum_{\zeta(\rho) = 0} \frac{x^\rho}{\rho} - \ln(2\pi)$$
$$|\rho| < 400$$



Approximations by Dirichlet series

Approximations by Dirichlet series

$$\Delta_N(s) = 1^{-s} + \delta_{N,2} 2^{-s} + \cdots + \delta_{N,N} N^{-s}$$

$$N = 2M + 1$$

$$\Delta_N(\overline{\rho_M}) = \cdots = \Delta_N(\overline{\rho_1}) = 0 = \Delta_N(\rho_1) = \cdots = \Delta(\rho_M)$$

Approximations by Dirichlet series

$$\Delta_N(s) = 1^{-s} + \delta_{N,2} 2^{-s} + \cdots + \delta_{N,N} N^{-s}$$

$$N = 2M + 1$$

$$\Delta_N(\overline{\rho_M}) = \cdots = \Delta_N(\overline{\rho_1}) = 0 = \Delta_N(\rho_1) = \cdots = \Delta(\rho_M)$$

Approximations by Dirichlet series

$$\Delta_N(s) = 1^{-s} + \delta_{N,2} 2^{-s} + \cdots + \delta_{N,N} N^{-s}$$

$$N = 2M + 1$$

$$\Delta_N(\overline{\rho_M}) = \cdots = \Delta_N(\overline{\rho_1}) = 0 = \Delta_N(\rho_1) = \cdots = \Delta(\rho_M)$$

$$\begin{vmatrix} 1 & 1 & \dots & 1 & 1 & 1 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ n^{-\overline{\rho_1}} & n^{-\rho_1} & \dots & n^{-\overline{\rho_M}} & n^{-\rho_M} & n^{-s} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ N^{-\overline{\rho_1}} & N^{-\rho_1} & \dots & N^{-\overline{\rho_M}} & N^{-\rho_M} & N^{-s} \end{vmatrix}$$

Approximations by Dirichlet series

$$\Delta_N(s) = 1^{-s} + \delta_{N,2} 2^{-s} + \cdots + \delta_{N,N} N^{-s}$$

$$N = 2M + 1$$

$$\Delta_N(\overline{\rho_M}) = \cdots = \Delta_N(\overline{\rho_1}) = 0 = \Delta_N(\rho_1) = \cdots = \Delta(\rho_M)$$

$$\begin{vmatrix} 1 & 1 & \dots & 1 & 1 & 1 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ n^{-\overline{\rho_1}} & n^{-\rho_1} & \dots & n^{-\overline{\rho_M}} & n^{-\rho_M} & n^{-s} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ N^{-\overline{\rho_1}} & N^{-\rho_1} & \dots & N^{-\overline{\rho_M}} & N^{-\rho_M} & N^{-s} \end{vmatrix} = \sum_{n=1}^N \tilde{\delta}_{N,n} n^{-s}$$

Approximations by Dirichlet series

$$\Delta_N(s) = 1^{-s} + \delta_{N,2} 2^{-s} + \cdots + \delta_{N,N} N^{-s}$$

$$N = 2M + 1$$

$$\Delta_N(\overline{\rho_M}) = \cdots = \Delta_N(\overline{\rho_1}) = 0 = \Delta_N(\rho_1) = \cdots = \Delta(\rho_M)$$

$$\begin{vmatrix} 1 & 1 & \dots & 1 & 1 & 1 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ n^{-\overline{\rho_1}} & n^{-\rho_1} & \dots & n^{-\overline{\rho_M}} & n^{-\rho_M} & n^{-s} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ N^{-\overline{\rho_1}} & N^{-\rho_1} & \dots & N^{-\overline{\rho_M}} & N^{-\rho_M} & N^{-s} \end{vmatrix} = \sum_{n=1}^N \tilde{\delta}_{N,n} n^{-s}$$

Approximations by Dirichlet series

$$\begin{vmatrix} 1 & 1 & \dots & 1 & 1 & 1 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ n^{-\rho_1} & n^{-\rho_1} & \dots & n^{-\rho_M} & n^{-\rho_M} & n^{-s} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ N^{-\rho_1} & N^{-\rho_1} & \dots & N^{-\rho_M} & N^{-\rho_M} & N^{-s} \end{vmatrix} = \sum_{n=1}^N \tilde{\delta}_{N,n} n^{-s}$$

Approximations by Dirichlet series

$$\begin{vmatrix} 1 & 1 & \dots & 1 & 1 & 1 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ n^{-\overline{\rho_1}} & n^{-\rho_1} & \dots & n^{-\overline{\rho_M}} & n^{-\rho_M} & n^{-s} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ N^{-\overline{\rho_1}} & N^{-\rho_1} & \dots & N^{-\overline{\rho_M}} & N^{-\rho_M} & N^{-s} \end{vmatrix} = \sum_{n=1}^N \tilde{\delta}_{N,n} n^{-s}$$

$$\tilde{\delta}_{N,n} = (-1)^{n+1} \begin{vmatrix} 1 & 1 & \dots & 1 & 1 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ (n-1)^{-\overline{\rho_1}} & (n-1)^{-\rho_1} & \dots & (n-1)^{-\overline{\rho_M}} & (n-1)^{-\rho_M} \\ (n+1)^{-\overline{\rho_1}} & (n+1)^{-\rho_1} & \dots & (n+1)^{-\overline{\rho_M}} & (n+1)^{-\rho_M} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ N^{-\overline{\rho_1}} & N^{-\rho_1} & \dots & N^{-\overline{\rho_M}} & N^{-\rho_M} \end{vmatrix}$$

Approximations by Dirichlet series

$$\begin{vmatrix} 1 & 1 & \dots & 1 & 1 & 1 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ n^{-\bar{\rho}_1} & n^{-\rho_1} & \dots & n^{-\bar{\rho}_M} & n^{-\rho_M} & n^{-s} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ N^{-\bar{\rho}_1} & N^{-\rho_1} & \dots & N^{-\bar{\rho}_M} & N^{-\rho_M} & N^{-s} \end{vmatrix} = \sum_{n=1}^N \tilde{\delta}_{N,n} n^{-s}$$

$$\tilde{\delta}_{N,n} = (-1)^{n+1} \begin{vmatrix} 1 & 1 & \dots & 1 & 1 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ (n-1)^{-\bar{\rho}_1} & (n-1)^{-\rho_1} & \dots & (n-1)^{-\bar{\rho}_M} & (n-1)^{-\rho_M} \\ (n+1)^{-\bar{\rho}_1} & (n+1)^{-\rho_1} & \dots & (n+1)^{-\bar{\rho}_M} & (n+1)^{-\rho_M} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ N^{-\bar{\rho}_1} & N^{-\rho_1} & \dots & N^{-\bar{\rho}_M} & N^{-\rho_M} \end{vmatrix}$$

$$\delta_{N,n} = \frac{\tilde{\delta}_{N,n}}{\tilde{\delta}_{N,1}}$$

Approximations by Dirichlet series

$$\begin{vmatrix} 1 & 1 & \dots & 1 & 1 & 1 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ n^{-\bar{\rho}_1} & n^{-\rho_1} & \dots & n^{-\bar{\rho}_M} & n^{-\rho_M} & n^{-s} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ N^{-\bar{\rho}_1} & N^{-\rho_1} & \dots & N^{-\bar{\rho}_M} & N^{-\rho_M} & N^{-s} \end{vmatrix} = \sum_{n=1}^N \tilde{\delta}_{N,n} n^{-s}$$

$$\tilde{\delta}_{N,n} = (-1)^{n+1} \begin{vmatrix} 1 & 1 & \dots & 1 & 1 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ (n-1)^{-\bar{\rho}_1} & (n-1)^{-\rho_1} & \dots & (n-1)^{-\bar{\rho}_M} & (n-1)^{-\rho_M} \\ (n+1)^{-\bar{\rho}_1} & (n+1)^{-\rho_1} & \dots & (n+1)^{-\bar{\rho}_M} & (n+1)^{-\rho_M} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ N^{-\bar{\rho}_1} & N^{-\rho_1} & \dots & N^{-\bar{\rho}_M} & N^{-\rho_M} \end{vmatrix}$$

$$\delta_{N,n} = \frac{\tilde{\delta}_{N,n}}{\tilde{\delta}_{N,1}} \quad \Delta_N(s) = \sum_{n=1}^N \delta_{N,n} n^{-s}$$

Trivial zeroes for $M = 1550$, $N = 2M + 1 = 3101$

Trivial zeroes for $M = 1550$, $N = 2M + 1 = 3101$

$$0 = \Delta_N(-2 - 1.884\ldots \cdot 10^{-1510})$$

Trivial zeroes for $M = 1550$, $N = 2M + 1 = 3101$

$$0 = \Delta_N(-2 - 1.884\ldots \cdot 10^{-1510})$$

$$0 = \Delta_N(-4 + 2.013\ldots \cdot 10^{-1504})$$

Trivial zeroes for $M = 1550$, $N = 2M + 1 = 3101$

$$0 = \Delta_N(-2 - 1.884 \dots \cdot 10^{-1510})$$

$$0 = \Delta_N(-4 + 2.013 \dots \cdot 10^{-1504})$$

$$0 = \Delta_N(-6 - 1.158 \dots \cdot 10^{-1498})$$

Trivial zeroes for $M = 1550$, $N = 2M + 1 = 3101$

$$0 = \Delta_N(-2 - 1.884 \dots \cdot 10^{-1510})$$

$$0 = \Delta_N(-4 + 2.013 \dots \cdot 10^{-1504})$$

$$0 = \Delta_N(-6 - 1.158 \dots \cdot 10^{-1498})$$

$$0 = \Delta_N(-8 + 4.508 \dots \cdot 10^{-1493})$$

Trivial zeroes for $M = 1550$, $N = 2M + 1 = 3101$

$$0 = \Delta_N(-2 - 1.884 \dots \cdot 10^{-1510})$$

$$0 = \Delta_N(-4 + 2.013 \dots \cdot 10^{-1504})$$

$$0 = \Delta_N(-6 - 1.158 \dots \cdot 10^{-1498})$$

$$0 = \Delta_N(-8 + 4.508 \dots \cdot 10^{-1493})$$

$$0 = \Delta_N(-10 - 1.316 \dots \cdot 10^{-1487})$$

Trivial zeroes for $M = 1550$, $N = 2M + 1 = 3101$

$$0 = \Delta_N(-2 - 1.884 \dots \cdot 10^{-1510})$$

$$0 = \Delta_N(-4 + 2.013 \dots \cdot 10^{-1504})$$

$$0 = \Delta_N(-6 - 1.158 \dots \cdot 10^{-1498})$$

$$0 = \Delta_N(-8 + 4.508 \dots \cdot 10^{-1493})$$

$$0 = \Delta_N(-10 - 1.316 \dots \cdot 10^{-1487})$$

$$0 = \Delta_N(-12 + 3.066 \dots \cdot 10^{-1482})$$

Trivial zeroes for $M = 1550$, $N = 2M + 1 = 3101$

$$0 = \Delta_N(-2 - 1.884 \dots \cdot 10^{-1510})$$

$$0 = \Delta_N(-4 + 2.013 \dots \cdot 10^{-1504})$$

$$0 = \Delta_N(-6 - 1.158 \dots \cdot 10^{-1498})$$

$$0 = \Delta_N(-8 + 4.508 \dots \cdot 10^{-1493})$$

$$0 = \Delta_N(-10 - 1.316 \dots \cdot 10^{-1487})$$

$$0 = \Delta_N(-12 + 3.066 \dots \cdot 10^{-1482})$$

$$0 = \Delta_N(-14 - 5.931 \dots \cdot 10^{-1477})$$

Trivial zeroes for $M = 1550$, $N = 2M + 1 = 3101$

$$0 = \Delta_N(-2 - 1.884 \dots \cdot 10^{-1510})$$

$$0 = \Delta_N(-4 + 2.013 \dots \cdot 10^{-1504})$$

$$0 = \Delta_N(-6 - 1.158 \dots \cdot 10^{-1498})$$

$$0 = \Delta_N(-8 + 4.508 \dots \cdot 10^{-1493})$$

$$0 = \Delta_N(-10 - 1.316 \dots \cdot 10^{-1487})$$

$$0 = \Delta_N(-12 + 3.066 \dots \cdot 10^{-1482})$$

$$0 = \Delta_N(-14 - 5.931 \dots \cdot 10^{-1477})$$

$$0 = \Delta_N(-16 + 9.796 \dots \cdot 10^{-1472})$$

Trivial zeroes for $M = 1550$, $N = 2M + 1 = 3101$

$$0 = \Delta_N(-2 - 1.884 \dots \cdot 10^{-1510})$$

$$0 = \Delta_N(-4 + 2.013 \dots \cdot 10^{-1504})$$

$$0 = \Delta_N(-6 - 1.158 \dots \cdot 10^{-1498})$$

$$0 = \Delta_N(-8 + 4.508 \dots \cdot 10^{-1493})$$

$$0 = \Delta_N(-10 - 1.316 \dots \cdot 10^{-1487})$$

$$0 = \Delta_N(-12 + 3.066 \dots \cdot 10^{-1482})$$

$$0 = \Delta_N(-14 - 5.931 \dots \cdot 10^{-1477})$$

$$0 = \Delta_N(-16 + 9.796 \dots \cdot 10^{-1472})$$

$$0 = \Delta_N(-18 - 1.410 \dots \cdot 10^{-1466})$$

Trivial zeroes for $M = 1550$, $N = 2M + 1 = 3101$

$$0 = \Delta_N(-2 - 1.884 \dots \cdot 10^{-1510})$$

$$0 = \Delta_N(-4 + 2.013 \dots \cdot 10^{-1504})$$

$$0 = \Delta_N(-6 - 1.158 \dots \cdot 10^{-1498})$$

$$0 = \Delta_N(-8 + 4.508 \dots \cdot 10^{-1493})$$

$$0 = \Delta_N(-10 - 1.316 \dots \cdot 10^{-1487})$$

$$0 = \Delta_N(-12 + 3.066 \dots \cdot 10^{-1482})$$

$$0 = \Delta_N(-14 - 5.931 \dots \cdot 10^{-1477})$$

$$0 = \Delta_N(-16 + 9.796 \dots \cdot 10^{-1472})$$

$$0 = \Delta_N(-18 - 1.410 \dots \cdot 10^{-1466})$$

$$0 = \Delta_N(-20 + 1.797 \dots \cdot 10^{-1461})$$

Trivial zeroes for $M = 1550$, $N = 2M + 1 = 3101$

$$0 = \Delta_N(-2 - 1.884 \dots \cdot 10^{-1510})$$

$$0 = \Delta_N(-4 + 2.013 \dots \cdot 10^{-1504})$$

$$0 = \Delta_N(-6 - 1.158 \dots \cdot 10^{-1498})$$

$$0 = \Delta_N(-8 + 4.508 \dots \cdot 10^{-1493})$$

$$0 = \Delta_N(-10 - 1.316 \dots \cdot 10^{-1487})$$

$$0 = \Delta_N(-12 + 3.066 \dots \cdot 10^{-1482})$$

$$0 = \Delta_N(-14 - 5.931 \dots \cdot 10^{-1477})$$

$$0 = \Delta_N(-16 + 9.796 \dots \cdot 10^{-1472})$$

$$0 = \Delta_N(-18 - 1.410 \dots \cdot 10^{-1466})$$

$$0 = \Delta_N(-20 + 1.797 \dots \cdot 10^{-1461})$$

$$0 = \Delta_N(-22 - 2.054 \dots \cdot 10^{-1456})$$

Trivial zeroes for $M = 1550$, $N = 2M + 1 = 3101$

$$0 = \Delta_N(-2 - 1.884 \dots \cdot 10^{-1510})$$

$$0 = \Delta_N(-4 + 2.013 \dots \cdot 10^{-1504})$$

$$0 = \Delta_N(-6 - 1.158 \dots \cdot 10^{-1498})$$

$$0 = \Delta_N(-8 + 4.508 \dots \cdot 10^{-1493})$$

$$0 = \Delta_N(-10 - 1.316 \dots \cdot 10^{-1487})$$

$$0 = \Delta_N(-12 + 3.066 \dots \cdot 10^{-1482})$$

$$0 = \Delta_N(-14 - 5.931 \dots \cdot 10^{-1477})$$

$$0 = \Delta_N(-16 + 9.796 \dots \cdot 10^{-1472})$$

$$0 = \Delta_N(-18 - 1.410 \dots \cdot 10^{-1466})$$

$$0 = \Delta_N(-20 + 1.797 \dots \cdot 10^{-1461})$$

$$0 = \Delta_N(-22 - 2.054 \dots \cdot 10^{-1456})$$

$$0 = \Delta_N(-24 + 2.126 \dots \cdot 10^{-1451})$$

Non-trivial zeroes for $M = 1550$, $N = 2M + 1 = 3101$

Non-trivial zeroes for $M = 1550$, $N = 2M + 1 = 3101$

$$0 = \Delta_N(\rho_{M+1} - 5.154\dots \cdot 10^{-1157} + 1.120\dots \cdot 10^{-1156}i)$$

Non-trivial zeroes for $M = 1550$, $N = 2M + 1 = 3101$

$$0 = \Delta_N(\rho_{M+1} - 5.154\dots \cdot 10^{-1157} + 1.120\dots \cdot 10^{-1156}i)$$

$$0 = \Delta_N(\rho_{M+201} - 4.922\dots \cdot 10^{-890} - 9.995\dots \cdot 10^{-891}i)$$

Non-trivial zeroes for $M = 1550$, $N = 2M + 1 = 3101$

$$0 = \Delta_N(\rho_{M+1} - 5.154 \dots \cdot 10^{-1157} + 1.120 \dots \cdot 10^{-1156}i)$$

$$0 = \Delta_N(\rho_{M+201} - 4.922 \dots \cdot 10^{-890} - 9.995 \dots \cdot 10^{-891}i)$$

$$0 = \Delta_N(\rho_{M+401} - 3.159 \dots \cdot 10^{-735} - 2.750 \dots \cdot 10^{-735}i)$$

Non-trivial zeroes for $M = 1550$, $N = 2M + 1 = 3101$

$$0 = \Delta_N(\rho_{M+1} - 5.154 \dots \cdot 10^{-1157} + 1.120 \dots \cdot 10^{-1156}i)$$

$$0 = \Delta_N(\rho_{M+201} - 4.922 \dots \cdot 10^{-890} - 9.995 \dots \cdot 10^{-891}i)$$

$$0 = \Delta_N(\rho_{M+401} - 3.159 \dots \cdot 10^{-735} - 2.750 \dots \cdot 10^{-735}i)$$

$$0 = \Delta_N(\rho_{M+601} + 8.765 \dots \cdot 10^{-619} + 4.575 \dots \cdot 10^{-618}i)$$

Non-trivial zeroes for $M = 1550$, $N = 2M + 1 = 3101$

$$0 = \Delta_N(\rho_{M+1} - 5.154 \dots \cdot 10^{-1157} + 1.120 \dots \cdot 10^{-1156}i)$$

$$0 = \Delta_N(\rho_{M+201} - 4.922 \dots \cdot 10^{-890} - 9.995 \dots \cdot 10^{-891}i)$$

$$0 = \Delta_N(\rho_{M+401} - 3.159 \dots \cdot 10^{-735} - 2.750 \dots \cdot 10^{-735}i)$$

$$0 = \Delta_N(\rho_{M+601} + 8.765 \dots \cdot 10^{-619} + 4.575 \dots \cdot 10^{-618}i)$$

$$0 = \Delta_N(\rho_{M+801} + 2.075 \dots \cdot 10^{-524} + 1.197 \dots \cdot 10^{-524}i)$$

Non-trivial zeroes for $M = 1550$, $N = 2M + 1 = 3101$

$$0 = \Delta_N(\rho_{M+1} - 5.154 \dots \cdot 10^{-1157} + 1.120 \dots \cdot 10^{-1156}i)$$

$$0 = \Delta_N(\rho_{M+201} - 4.922 \dots \cdot 10^{-890} - 9.995 \dots \cdot 10^{-891}i)$$

$$0 = \Delta_N(\rho_{M+401} - 3.159 \dots \cdot 10^{-735} - 2.750 \dots \cdot 10^{-735}i)$$

$$0 = \Delta_N(\rho_{M+601} + 8.765 \dots \cdot 10^{-619} + 4.575 \dots \cdot 10^{-618}i)$$

$$0 = \Delta_N(\rho_{M+801} + 2.075 \dots \cdot 10^{-524} + 1.197 \dots \cdot 10^{-524}i)$$

$$0 = \Delta_N(\rho_{M+1001} + 1.980 \dots \cdot 10^{-447} - 3.397 \dots \cdot 10^{-448}i)$$

Non-trivial zeroes for $M = 1550$, $N = 2M + 1 = 3101$

$$0 = \Delta_N(\rho_{M+1} - 5.154 \dots \cdot 10^{-1157} + 1.120 \dots \cdot 10^{-1156}i)$$

$$0 = \Delta_N(\rho_{M+201} - 4.922 \dots \cdot 10^{-890} - 9.995 \dots \cdot 10^{-891}i)$$

$$0 = \Delta_N(\rho_{M+401} - 3.159 \dots \cdot 10^{-735} - 2.750 \dots \cdot 10^{-735}i)$$

$$0 = \Delta_N(\rho_{M+601} + 8.765 \dots \cdot 10^{-619} + 4.575 \dots \cdot 10^{-618}i)$$

$$0 = \Delta_N(\rho_{M+801} + 2.075 \dots \cdot 10^{-524} + 1.197 \dots \cdot 10^{-524}i)$$

$$0 = \Delta_N(\rho_{M+1001} + 1.980 \dots \cdot 10^{-447} - 3.397 \dots \cdot 10^{-448}i)$$

$$0 = \Delta_N(\rho_{M+1201} - 1.034 \dots \cdot 10^{-381} - 1.354 \dots \cdot 10^{-382}i)$$

Non-trivial zeroes for $M = 1550$, $N = 2M + 1 = 3101$

$$0 = \Delta_N(\rho_{M+1} - 5.154 \dots \cdot 10^{-1157} + 1.120 \dots \cdot 10^{-1156}i)$$

$$0 = \Delta_N(\rho_{M+201} - 4.922 \dots \cdot 10^{-890} - 9.995 \dots \cdot 10^{-891}i)$$

$$0 = \Delta_N(\rho_{M+401} - 3.159 \dots \cdot 10^{-735} - 2.750 \dots \cdot 10^{-735}i)$$

$$0 = \Delta_N(\rho_{M+601} + 8.765 \dots \cdot 10^{-619} + 4.575 \dots \cdot 10^{-618}i)$$

$$0 = \Delta_N(\rho_{M+801} + 2.075 \dots \cdot 10^{-524} + 1.197 \dots \cdot 10^{-524}i)$$

$$0 = \Delta_N(\rho_{M+1001} + 1.980 \dots \cdot 10^{-447} - 3.397 \dots \cdot 10^{-448}i)$$

$$0 = \Delta_N(\rho_{M+1201} - 1.034 \dots \cdot 10^{-381} - 1.354 \dots \cdot 10^{-382}i)$$

$$0 = \Delta_N(\rho_{M+1401} + 1.466 \dots \cdot 10^{-326} - 1.835 \dots \cdot 10^{-326}i)$$

Non-trivial zeroes for $M = 1550$, $N = 2M + 1 = 3101$

$$0 = \Delta_N(\rho_{M+1} - 5.154 \dots \cdot 10^{-1157} + 1.120 \dots \cdot 10^{-1156}i)$$

$$0 = \Delta_N(\rho_{M+201} - 4.922 \dots \cdot 10^{-890} - 9.995 \dots \cdot 10^{-891}i)$$

$$0 = \Delta_N(\rho_{M+401} - 3.159 \dots \cdot 10^{-735} - 2.750 \dots \cdot 10^{-735}i)$$

$$0 = \Delta_N(\rho_{M+601} + 8.765 \dots \cdot 10^{-619} + 4.575 \dots \cdot 10^{-618}i)$$

$$0 = \Delta_N(\rho_{M+801} + 2.075 \dots \cdot 10^{-524} + 1.197 \dots \cdot 10^{-524}i)$$

$$0 = \Delta_N(\rho_{M+1001} + 1.980 \dots \cdot 10^{-447} - 3.397 \dots \cdot 10^{-448}i)$$

$$0 = \Delta_N(\rho_{M+1201} - 1.034 \dots \cdot 10^{-381} - 1.354 \dots \cdot 10^{-382}i)$$

$$0 = \Delta_N(\rho_{M+1401} + 1.466 \dots \cdot 10^{-326} - 1.835 \dots \cdot 10^{-326}i)$$

$$0 = \Delta_N(\rho_{M+1601} + 2.281 \dots \cdot 10^{-278} - 3.603 \dots \cdot 10^{-278}i)$$

Non-trivial zeroes for $M = 1550$, $N = 2M + 1 = 3101$

$$0 = \Delta_N(\rho_{M+1} - 5.154 \dots \cdot 10^{-1157} + 1.120 \dots \cdot 10^{-1156}i)$$

$$0 = \Delta_N(\rho_{M+201} - 4.922 \dots \cdot 10^{-890} - 9.995 \dots \cdot 10^{-891}i)$$

$$0 = \Delta_N(\rho_{M+401} - 3.159 \dots \cdot 10^{-735} - 2.750 \dots \cdot 10^{-735}i)$$

$$0 = \Delta_N(\rho_{M+601} + 8.765 \dots \cdot 10^{-619} + 4.575 \dots \cdot 10^{-618}i)$$

$$0 = \Delta_N(\rho_{M+801} + 2.075 \dots \cdot 10^{-524} + 1.197 \dots \cdot 10^{-524}i)$$

$$0 = \Delta_N(\rho_{M+1001} + 1.980 \dots \cdot 10^{-447} - 3.397 \dots \cdot 10^{-448}i)$$

$$0 = \Delta_N(\rho_{M+1201} - 1.034 \dots \cdot 10^{-381} - 1.354 \dots \cdot 10^{-382}i)$$

$$0 = \Delta_N(\rho_{M+1401} + 1.466 \dots \cdot 10^{-326} - 1.835 \dots \cdot 10^{-326}i)$$

$$0 = \Delta_N(\rho_{M+1601} + 2.281 \dots \cdot 10^{-278} - 3.603 \dots \cdot 10^{-278}i)$$

$$0 = \Delta_N(\rho_{M+1801} - 7.799 \dots \cdot 10^{-237} - 3.726 \dots \cdot 10^{-237}i)$$

Non-trivial zeroes for $M = 1550$, $N = 2M + 1 = 3101$

$$\begin{aligned}0 &= \Delta_N(\rho_{M+1} - 5.154 \dots \cdot 10^{-1157} + 1.120 \dots \cdot 10^{-1156}i) \\0 &= \Delta_N(\rho_{M+201} - 4.922 \dots \cdot 10^{-890} - 9.995 \dots \cdot 10^{-891}i) \\0 &= \Delta_N(\rho_{M+401} - 3.159 \dots \cdot 10^{-735} - 2.750 \dots \cdot 10^{-735}i) \\0 &= \Delta_N(\rho_{M+601} + 8.765 \dots \cdot 10^{-619} + 4.575 \dots \cdot 10^{-618}i) \\0 &= \Delta_N(\rho_{M+801} + 2.075 \dots \cdot 10^{-524} + 1.197 \dots \cdot 10^{-524}i) \\0 &= \Delta_N(\rho_{M+1001} + 1.980 \dots \cdot 10^{-447} - 3.397 \dots \cdot 10^{-448}i) \\0 &= \Delta_N(\rho_{M+1201} - 1.034 \dots \cdot 10^{-381} - 1.354 \dots \cdot 10^{-382}i) \\0 &= \Delta_N(\rho_{M+1401} + 1.466 \dots \cdot 10^{-326} - 1.835 \dots \cdot 10^{-326}i) \\0 &= \Delta_N(\rho_{M+1601} + 2.281 \dots \cdot 10^{-278} - 3.603 \dots \cdot 10^{-278}i) \\0 &= \Delta_N(\rho_{M+1801} - 7.799 \dots \cdot 10^{-237} - 3.726 \dots \cdot 10^{-237}i) \\0 &= \Delta_N(\rho_{M+2001} + 5.921 \dots \cdot 10^{-201} - 6.855 \dots \cdot 10^{-201}i)\end{aligned}$$

Non-trivial zeroes for $M = 1550$, $N = 2M + 1 = 3101$

$$\begin{aligned} 0 &= \Delta_N(\rho_{M+1} - 5.154 \dots \cdot 10^{-1157} + 1.120 \dots \cdot 10^{-1156}i) \\ 0 &= \Delta_N(\rho_{M+201} - 4.922 \dots \cdot 10^{-890} - 9.995 \dots \cdot 10^{-891}i) \\ 0 &= \Delta_N(\rho_{M+401} - 3.159 \dots \cdot 10^{-735} - 2.750 \dots \cdot 10^{-735}i) \\ 0 &= \Delta_N(\rho_{M+601} + 8.765 \dots \cdot 10^{-619} + 4.575 \dots \cdot 10^{-618}i) \\ 0 &= \Delta_N(\rho_{M+801} + 2.075 \dots \cdot 10^{-524} + 1.197 \dots \cdot 10^{-524}i) \\ 0 &= \Delta_N(\rho_{M+1001} + 1.980 \dots \cdot 10^{-447} - 3.397 \dots \cdot 10^{-448}i) \\ 0 &= \Delta_N(\rho_{M+1201} - 1.034 \dots \cdot 10^{-381} - 1.354 \dots \cdot 10^{-382}i) \\ 0 &= \Delta_N(\rho_{M+1401} + 1.466 \dots \cdot 10^{-326} - 1.835 \dots \cdot 10^{-326}i) \\ 0 &= \Delta_N(\rho_{M+1601} + 2.281 \dots \cdot 10^{-278} - 3.603 \dots \cdot 10^{-278}i) \\ 0 &= \Delta_N(\rho_{M+1801} - 7.799 \dots \cdot 10^{-237} - 3.726 \dots \cdot 10^{-237}i) \\ 0 &= \Delta_N(\rho_{M+2001} + 5.921 \dots \cdot 10^{-201} - 6.855 \dots \cdot 10^{-201}i) \\ 0 &= \Delta_N(\rho_{M+2201} + 8.049 \dots \cdot 10^{-170} + 1.359 \dots \cdot 10^{-169}i) \end{aligned}$$

Non-trivial zeroes for $M = 1550$, $N = 2M + 1 = 3101$

$$\begin{aligned} 0 &= \Delta_N(\rho_{M+1} - 5.154 \dots \cdot 10^{-1157} + 1.120 \dots \cdot 10^{-1156}i) \\ 0 &= \Delta_N(\rho_{M+201} - 4.922 \dots \cdot 10^{-890} - 9.995 \dots \cdot 10^{-891}i) \\ 0 &= \Delta_N(\rho_{M+401} - 3.159 \dots \cdot 10^{-735} - 2.750 \dots \cdot 10^{-735}i) \\ 0 &= \Delta_N(\rho_{M+601} + 8.765 \dots \cdot 10^{-619} + 4.575 \dots \cdot 10^{-618}i) \\ 0 &= \Delta_N(\rho_{M+801} + 2.075 \dots \cdot 10^{-524} + 1.197 \dots \cdot 10^{-524}i) \\ 0 &= \Delta_N(\rho_{M+1001} + 1.980 \dots \cdot 10^{-447} - 3.397 \dots \cdot 10^{-448}i) \\ 0 &= \Delta_N(\rho_{M+1201} - 1.034 \dots \cdot 10^{-381} - 1.354 \dots \cdot 10^{-382}i) \\ 0 &= \Delta_N(\rho_{M+1401} + 1.466 \dots \cdot 10^{-326} - 1.835 \dots \cdot 10^{-326}i) \\ 0 &= \Delta_N(\rho_{M+1601} + 2.281 \dots \cdot 10^{-278} - 3.603 \dots \cdot 10^{-278}i) \\ 0 &= \Delta_N(\rho_{M+1801} - 7.799 \dots \cdot 10^{-237} - 3.726 \dots \cdot 10^{-237}i) \\ 0 &= \Delta_N(\rho_{M+2001} + 5.921 \dots \cdot 10^{-201} - 6.855 \dots \cdot 10^{-201}i) \\ 0 &= \Delta_N(\rho_{M+2201} + 8.049 \dots \cdot 10^{-170} + 1.359 \dots \cdot 10^{-169}i) \\ 0 &= \Delta_N(\rho_{M+2401} - 2.001 \dots \cdot 10^{-142} - 7.023 \dots \cdot 10^{-142}i) \end{aligned}$$

Extra zeroes for $M = 1550$, $N = 2M + 1 = 3101$

$$\Delta_N(s) \doteq \eta(s) = 1^{-s} - 2^{-s} + 3^{-s} - 4^{-s} + \dots = (1 - 2 \cdot 2^{-s})\zeta(s)$$

Extra zeroes for $M = 1550$, $N = 2M + 1 = 3101$

$$\Delta_N(s) = \eta(s) = 1^{-s} - 2^{-s} + 3^{-s} - 4^{-s} + \dots = (1 - 2 \cdot 2^{-s})\zeta(s)$$

$$1 - 2 \cdot 2^{-s} = 0 \iff s = s_k = 1 + \frac{2\pi k}{\ln(2)}i, \quad k = 0, \pm 1, \pm 2, \dots$$

$$0 = \Delta_N(s_{50} - 5.481\dots \cdot 10^{-133} - 5.546\dots \cdot 10^{-133}i)$$

Extra zeroes for $M = 1550$, $N = 2M + 1 = 3101$

$$\Delta_N(s) = \eta(s) = 1^{-s} - 2^{-s} + 3^{-s} - 4^{-s} + \dots = (1 - 2 \cdot 2^{-s})\zeta(s)$$

$$1 - 2 \cdot 2^{-s} = 0 \iff s = s_k = 1 + \frac{2\pi k}{\ln(2)}i, \quad k = 0, \pm 1, \pm 2, \dots$$

$$0 = \Delta_N(s_{50} - 5.481\dots \cdot 10^{-133} - 5.546\dots \cdot 10^{-133}i)$$

$$0 = \Delta_N(s_{100} - 1.109\dots \cdot 10^{-132} - 1.306\dots \cdot 10^{-134}i)$$

Extra zeroes for $M = 1550$, $N = 2M + 1 = 3101$

$$\Delta_N(s) = \eta(s) = 1^{-s} - 2^{-s} + 3^{-s} - 4^{-s} + \dots = (1 - 2 \cdot 2^{-s})\zeta(s)$$

$$1 - 2 \cdot 2^{-s} = 0 \iff s = s_k = 1 + \frac{2\pi k}{\ln(2)}i, \quad k = 0, \pm 1, \pm 2, \dots$$

$$0 = \Delta_N(s_{50} - 5.481\dots \cdot 10^{-133} - 5.546\dots \cdot 10^{-133}i)$$

$$0 = \Delta_N(s_{100} - 1.109\dots \cdot 10^{-132} - 1.306\dots \cdot 10^{-134}i)$$

$$0 = \Delta_N(s_{150} - 5.743\dots \cdot 10^{-133} + 5.543\dots \cdot 10^{-133}i)$$

Extra zeroes for $M = 1550$, $N = 2M + 1 = 3101$

$$\Delta_N(s) = \eta(s) = 1^{-s} - 2^{-s} + 3^{-s} - 4^{-s} + \dots = (1 - 2 \cdot 2^{-s})\zeta(s)$$

$$1 - 2 \cdot 2^{-s} = 0 \iff s = s_k = 1 + \frac{2\pi k}{\ln(2)}i, \quad k = 0, \pm 1, \pm 2, \dots$$

$$0 = \Delta_N(s_{50} - 5.481\dots \cdot 10^{-133} - 5.546\dots \cdot 10^{-133}i)$$

$$0 = \Delta_N(s_{100} - 1.109\dots \cdot 10^{-132} - 1.306\dots \cdot 10^{-134}i)$$

$$0 = \Delta_N(s_{150} - 5.743\dots \cdot 10^{-133} + 5.543\dots \cdot 10^{-133}i)$$

$$0 = \Delta_N(s_{200} - 6.157\dots \cdot 10^{-136} + 2.613\dots \cdot 10^{-134}i)$$

Extra zeroes for $M = 1550$, $N = 2M + 1 = 3101$

$$\Delta_N(s) = \eta(s) = 1^{-s} - 2^{-s} + 3^{-s} - 4^{-s} + \dots = (1 - 2 \cdot 2^{-s})\zeta(s)$$

$$1 - 2 \cdot 2^{-s} = 0 \iff s = s_k = 1 + \frac{2\pi k}{\ln(2)}i, \quad k = 0, \pm 1, \pm 2, \dots$$

$$0 = \Delta_N(s_{50} - 5.481\dots \cdot 10^{-133} - 5.546\dots \cdot 10^{-133}i)$$

$$0 = \Delta_N(s_{100} - 1.109\dots \cdot 10^{-132} - 1.306\dots \cdot 10^{-134}i)$$

$$0 = \Delta_N(s_{150} - 5.743\dots \cdot 10^{-133} + 5.543\dots \cdot 10^{-133}i)$$

$$0 = \Delta_N(s_{200} - 6.157\dots \cdot 10^{-136} + 2.613\dots \cdot 10^{-134}i)$$

$$0 = \Delta_N(s_{250} - 5.220\dots \cdot 10^{-133} - 5.537\dots \cdot 10^{-133}i)$$

Extra zeroes for $M = 1550$, $N = 2M + 1 = 3101$

$$\Delta_N(s) = \eta(s) = 1^{-s} - 2^{-s} + 3^{-s} - 4^{-s} + \dots = (1 - 2 \cdot 2^{-s})\zeta(s)$$

$$1 - 2 \cdot 2^{-s} = 0 \iff s = s_k = 1 + \frac{2\pi k}{\ln(2)}i, \quad k = 0, \pm 1, \pm 2, \dots$$

$$0 = \Delta_N(s_{50} - 5.481\dots \cdot 10^{-133} - 5.546\dots \cdot 10^{-133}i)$$

$$0 = \Delta_N(s_{100} - 1.109\dots \cdot 10^{-132} - 1.306\dots \cdot 10^{-134}i)$$

$$0 = \Delta_N(s_{150} - 5.743\dots \cdot 10^{-133} + 5.543\dots \cdot 10^{-133}i)$$

$$0 = \Delta_N(s_{200} - 6.157\dots \cdot 10^{-136} + 2.613\dots \cdot 10^{-134}i)$$

$$0 = \Delta_N(s_{250} - 5.220\dots \cdot 10^{-133} - 5.537\dots \cdot 10^{-133}i)$$

$$0 = \Delta_N(s_{300} - 1.108\dots \cdot 10^{-132} - 3.917\dots \cdot 10^{-134}i)$$

Extra zeroes for $M = 1550$, $N = 2M + 1 = 3101$

$$\Delta_N(s) \doteq \eta(s) = 1^{-s} - 2^{-s} + 3^{-s} - 4^{-s} + \dots = (1 - 2 \cdot 2^{-s})\zeta(s)$$

$$1 - 2 \cdot 2^{-s} = 0 \iff s = s_k = 1 + \frac{2\pi k}{\ln(2)}i, \quad k = 0, \pm 1, \pm 2, \dots$$

$$0 = \Delta_N(s_{50} - 5.481\dots \cdot 10^{-133} - 5.546\dots \cdot 10^{-133}i)$$

$$0 = \Delta_N(s_{100} - 1.109\dots \cdot 10^{-132} - 1.306\dots \cdot 10^{-134}i)$$

$$0 = \Delta_N(s_{150} - 5.743\dots \cdot 10^{-133} + 5.543\dots \cdot 10^{-133}i)$$

$$0 = \Delta_N(s_{200} - 6.157\dots \cdot 10^{-136} + 2.613\dots \cdot 10^{-134}i)$$

$$0 = \Delta_N(s_{250} - 5.220\dots \cdot 10^{-133} - 5.537\dots \cdot 10^{-133}i)$$

$$0 = \Delta_N(s_{300} - 1.108\dots \cdot 10^{-132} - 3.917\dots \cdot 10^{-134}i)$$

$$0 = \Delta_N(s_{350} - 6.004\dots \cdot 10^{-133} + 5.528\dots \cdot 10^{-133}i)$$

Extra zeroes for $M = 1550$, $N = 2M + 1 = 3101$

$$\Delta_N(s) \doteq \eta(s) = 1^{-s} - 2^{-s} + 3^{-s} - 4^{-s} + \dots = (1 - 2 \cdot 2^{-s})\zeta(s)$$

$$1 - 2 \cdot 2^{-s} = 0 \iff s = s_k = 1 + \frac{2\pi k}{\ln(2)}i, \quad k = 0, \pm 1, \pm 2, \dots$$

$$0 = \Delta_N(s_{50} - 5.481\dots \cdot 10^{-133} - 5.546\dots \cdot 10^{-133}i)$$

$$0 = \Delta_N(s_{100} - 1.109\dots \cdot 10^{-132} - 1.306\dots \cdot 10^{-134}i)$$

$$0 = \Delta_N(s_{150} - 5.743\dots \cdot 10^{-133} + 5.543\dots \cdot 10^{-133}i)$$

$$0 = \Delta_N(s_{200} - 6.157\dots \cdot 10^{-136} + 2.613\dots \cdot 10^{-134}i)$$

$$0 = \Delta_N(s_{250} - 5.220\dots \cdot 10^{-133} - 5.537\dots \cdot 10^{-133}i)$$

$$0 = \Delta_N(s_{300} - 1.108\dots \cdot 10^{-132} - 3.917\dots \cdot 10^{-134}i)$$

$$0 = \Delta_N(s_{350} - 6.004\dots \cdot 10^{-133} + 5.528\dots \cdot 10^{-133}i)$$

$$0 = \Delta_N(s_{400} - 2.461\dots \cdot 10^{-135} + 5.220\dots \cdot 10^{-134}i)$$

Extra zeroes for $M = 1550$, $N = 2M + 1 = 3101$

$$\Delta_N(s) \doteq \eta(s) = 1^{-s} - 2^{-s} + 3^{-s} - 4^{-s} + \dots = (1 - 2 \cdot 2^{-s})\zeta(s)$$

$$1 - 2 \cdot 2^{-s} = 0 \iff s = s_k = 1 + \frac{2\pi k}{\ln(2)}i, \quad k = 0, \pm 1, \pm 2, \dots$$

$$0 = \Delta_N(s_{50} - 5.481\dots \cdot 10^{-133} - 5.546\dots \cdot 10^{-133}i)$$

$$0 = \Delta_N(s_{100} - 1.109\dots \cdot 10^{-132} - 1.306\dots \cdot 10^{-134}i)$$

$$0 = \Delta_N(s_{150} - 5.743\dots \cdot 10^{-133} + 5.543\dots \cdot 10^{-133}i)$$

$$0 = \Delta_N(s_{200} - 6.157\dots \cdot 10^{-136} + 2.613\dots \cdot 10^{-134}i)$$

$$0 = \Delta_N(s_{250} - 5.220\dots \cdot 10^{-133} - 5.537\dots \cdot 10^{-133}i)$$

$$0 = \Delta_N(s_{300} - 1.108\dots \cdot 10^{-132} - 3.917\dots \cdot 10^{-134}i)$$

$$0 = \Delta_N(s_{350} - 6.004\dots \cdot 10^{-133} + 5.528\dots \cdot 10^{-133}i)$$

$$0 = \Delta_N(s_{400} - 2.461\dots \cdot 10^{-135} + 5.220\dots \cdot 10^{-134}i)$$

Notation

Notation

$$\mu_{N,n} = \sum_{m|n} \mu\left(\frac{n}{m}\right) \delta_{N,m}$$

Notation

$$\mu_{N,n} = \sum_{m|n} \mu\left(\frac{n}{m}\right) \delta_{N,m}$$

$$\mu(k) = \begin{cases} (-1)^m, & \text{if } k \text{ is the product of } m \text{ different primes} \\ 0, & \text{otherwise} \end{cases}$$

Notation

$$\mu_{N,n} = \sum_{m|n} \mu\left(\frac{n}{m}\right) \delta_{N,m}$$

$$\mu(k) = \begin{cases} (-1)^m, & \text{if } k \text{ is the product of } m \text{ different primes} \\ 0, & \text{otherwise} \end{cases}$$

$$\mu_{N,1} = 1$$

$$\mu_{N,2} = \delta_{N,2} - 1$$

$$\mu_{N,3} = \delta_{N,3} - 1$$

$$\mu_{N,4} = \delta_{N,4} - \delta_{N,2}$$

$$\mu_{N,5} = \delta_{N,5} - 1$$

$$\mu_{N,6} = \delta_{N,6} - \delta_{N,3} - \delta_{N,2} + 1$$

$$\mu_{N,7} = \delta_{N,7} - 1$$

$$\mu_{N,8} = \delta_{N,8} - \delta_{N,4}$$

$$\mu_{N,9} = \delta_{N,9} - \delta_{N,3}$$

Case $N = 2001$


$$\begin{aligned}\mu_{2001,2} &= -2 + 9.93613 \dots \cdot 10^{-86} \\ \mu_{2001,3} &= -1.49042 \dots \cdot 10^{-85} \\ \mu_{2001,4} &= -1.35392 \dots \cdot 10^{-199} \\ \mu_{2001,5} &= +4.28708 \dots \cdot 10^{-297} \\ \mu_{2001,6} &= -1.39904 \dots \cdot 10^{-377} \\ \mu_{2001,7} &= -8.46908 \dots \cdot 10^{-444} \\ \mu_{2001,8} &= -3.00897 \dots \cdot 10^{-499} \\ \mu_{2001,9} &= +2.56119 \dots \cdot 10^{-546} \\ \mu_{2001,10} &= +9.47153 \dots \cdot 10^{-587} \\ \mu_{2001,11} &= -2.22088 \dots \cdot 10^{-622} \\ \mu_{2001,12} &= +1.65346 \dots \cdot 10^{-653} \\ \mu_{2001,13} &= -1.33219 \dots \cdot 10^{-680} \\ \mu_{2001,14} &= -2.89063 \dots \cdot 10^{-705} \\ \mu_{2001,15} &= -2.27283 \dots \cdot 10^{-726}\end{aligned}$$

Case $N = 2001$


$$\begin{aligned}\delta_{N,3} - \delta_{N,1} &= -1.49042 \dots \cdot 10^{-85} \\ \delta_{N,4} - \delta_{N,2} &= -1.35392 \dots \cdot 10^{-199} \\ \delta_{N,5} - \delta_{N,1} &= +4.28708 \dots \cdot 10^{-297} \\ \delta_{N,6} - \delta_{N,3} - \delta_{N,2} + \delta_{N,1} &= -1.39904 \dots \cdot 10^{-377} \\ \delta_{N,7} - \delta_{N,1} &= -8.46908 \dots \cdot 10^{-444} \\ \delta_{N,8} - \delta_{N,4} &= -3.00897 \dots \cdot 10^{-499} \\ \delta_{N,9} - \delta_{N,3} &= +2.56119 \dots \cdot 10^{-546} \\ \delta_{N,10} - \delta_{N,5} - \delta_{N,2} + \delta_{N,1} &= +9.47153 \dots \cdot 10^{-587} \\ \delta_{N,11} - \delta_{N,1} &= -2.22088 \dots \cdot 10^{-622} \\ \delta_{N,12} - \delta_{N,6} - \delta_{N,4} + \delta_{N,2} &= +1.65346 \dots \cdot 10^{-653} \\ \delta_{N,13} - \delta_{N,1} &= -1.33219 \dots \cdot 10^{-680} \\ \delta_{N,14} - \delta_{N,7} - \delta_{N,2} + \delta_{N,1} &= -2.89063 \dots \cdot 10^{-705} \\ \delta_{N,15} - \delta_{N,5} - \delta_{N,3} + \delta_{N,3} &= -2.27283 \dots \cdot 10^{-726}\end{aligned}$$

Coefficients $\delta_{3101,n}$, red for even n , blue for odd n

Coefficients $\delta_{3101,n}$, red for even n , blue for odd n



Coefficients $\delta_{3101,n}$, red for even n , blue for odd n



$$\delta_{3101,n}$$


$$\delta_{3101,n}-1$$

$$|\delta_{3101,n}-1|$$


$$\log_{10}|\delta_{3101,n}-1|$$

Plot of $\log_{10} |\delta_{3101,n} - 1|$


Plot of $\log_{10} |\delta_{3101,n} - 1|$




Plot of $\log_{10} |\delta_{3101,n} - 1|$




Plot of $\log_{10} |\delta_{3101,n} - 1|$




Plot of $\log_{10} |\delta_{3101,n} - 1|$




Plot of $\log_{10} |\delta_{3101,n} - 1|$




Plot of $\log_{10} |\delta_{3101,n} - 1|$




Plot of $\log_{10} |\delta_{3101,n} - 1|$




Plot of $\log_{10} |\delta_{3101,n} - 1|$




Plot of $\log_{10} |\delta_{3101,n} - 1|$ = Sieve of Eratosthenes



Plot of $\log_{10} |\delta_{3101,n} - 1|$




Scaled Plot of $\ln |\delta_{N,n} - (-1)^{n+1}|/N$ for $N = 6001$




$$\ell(a) = (a - 1) \ln(1 - a) - 2a \ln(a) + (a + 1) \ln(a + 1) - \ln(2\sqrt{2} + 3)$$

Plot of $\log_{10} |\delta_{6001,n} - 1|$ (repeated)




Finer Structure: Plot of $\log_{10} |\delta_{10001,3m} - \delta_{10001,3}|$


Finer Structure: Plot of $\log_{10} |\delta_{10001,3m} - \delta_{10001,3}|$




Finer Structure: Plot of $\log_{10} |\delta_{10001,3m} - \delta_{10001,3}|$




Finer Structure: Plot of $\log_{10} |\delta_{10001,3m} - \delta_{10001,3}|$




Finer Structure: Plot of $\log_{10} |\delta_{10001,3m} - \delta_{10001,3}|$




Finer Structure: Plot of $\log_{10} |\delta_{10001,3m} - \delta_{10001,3}|$



Finer Structure: Plot of $\log_{10} |\delta_{10001,3m} - \delta_{10001,3}|$




Finer Structure: Plot of $\log_{10} |\delta_{10001,3m} - \delta_{10001,3}|$



Finer Structure: Plot of $\log_{10} |\delta_{10001,3m} - \delta_{10001,3}|$

= Eratosthenes Sieve with primes order 2, 5, 7, 3, 11, 13, ...



Expected Fractal Structure

Let n range over the arithmetical progression $d, 2d, \dots, md, \dots$ with

$$d = 2^{k_2} 3^{k_3} 5^{k_5} \dots$$

Corresponding Eratosthenes sublevel splits according to the divisibility of m by p_1, p_2, \dots where these prime numbers are ordered in such a way that

$$p_1^{k_{p_1}+1} < p_2^{k_{p_2}+1} < \dots < p_j^{k_{p_j}+1} < \dots$$

Expected Fractal Structure

Let n range over the arithmetical progression $d, 2d, \dots, md, \dots$ with

$$d = 2^{k_2} 3^{k_3} 5^{k_5} \dots$$

Corresponding Eratosthenes sublevel splits according to the divisibility of m by p_1, p_2, \dots where these prime numbers are ordered in such a way that

$$p_1^{k_{p_1}+1} < p_2^{k_{p_2}+1} < \dots < p_j^{k_{p_j}+1} < \dots$$

In the previous example $m = 3$, hence $k_2 = 0, k_3 = 1, k_5 = k_7 = \dots = 0$ and $p_1 = 2, p_2 = 5, p_3 = 7, p_4 = 3, p_5 = 11, p_6 = 13, \dots$ according to

$$2^1 < 5^1 < 7^1 < 3^2 < 11^1 < 13^2 < \dots$$

Zeroes from Euler Product

$$\begin{aligned}\zeta(s) &= 1^{-s} + 2^{-s} + \cdots + k^{-s} + \dots \\ &= \prod_{p \text{ is prime}} \frac{1}{1 - p^{-s}}\end{aligned}$$

Zeroes from Euler Product

$$\begin{aligned}\zeta(s) &= 1^{-s} + 2^{-s} + \cdots + k^{-s} + \dots \\ &= \prod_{p \text{ is prime}} \frac{1}{1 - p^{-s}}\end{aligned}$$

$$\zeta_2(s) = \prod_{\substack{p \text{ is prime} \\ p \neq 2}} \frac{1}{1 - p^{-s}}$$

Zeroes from Euler Product

$$\begin{aligned}\zeta(s) &= 1^{-s} + 2^{-s} + \cdots + k^{-s} + \dots \\ &= \prod_{p \text{ is prime}} \frac{1}{1 - p^{-s}}\end{aligned}$$

$$\begin{aligned}\zeta_2(s) &= \prod_{\substack{p \text{ is prime} \\ p \neq 2}} \frac{1}{1 - p^{-s}} \\ &= 1^{-s} + 3^{-s} + \cdots + (2k+1)^{-s} + \dots\end{aligned}$$

Zeroes from Euler Product

$$\begin{aligned}\zeta(s) &= 1^{-s} + 2^{-s} + \cdots + k^{-s} + \dots \\ &= \prod_{p \text{ is prime}} \frac{1}{1 - p^{-s}}\end{aligned}$$

$$\begin{aligned}\zeta_2(s) &= \prod_{\substack{p \text{ is prime} \\ p \neq 2}} \frac{1}{1 - p^{-s}} \\ &= 1^{-s} + 3^{-s} + \cdots + (2k+1)^{-s} + \dots \\ &= L(2, \chi_1, s)\end{aligned}$$

Zeroes from Euler Product

$$\begin{aligned}\zeta(s) &= 1^{-s} + 2^{-s} + \cdots + k^{-s} + \dots \\ &= \prod_{p \text{ is prime}} \frac{1}{1 - p^{-s}}\end{aligned}$$

$$\begin{aligned}\zeta_2(s) &= \prod_{\substack{p \text{ is prime} \\ p \neq 2}} \frac{1}{1 - p^{-s}} \\ &= 1^{-s} + 3^{-s} + \cdots + (2k+1)^{-s} + \dots \\ &= L(2, \chi_1, s) \\ &= (1 - 2^{-s})\zeta(s)\end{aligned}$$

An Example

$$\zeta_2(s) = (1 - 2^{-s})\zeta(s)$$

An Example

$$\zeta_2(s) = (1 - 2^{-s})\zeta(s)$$

Let us take 100 (pairs of conjugate) zeros of the zeta function,

$$\frac{1}{2} \pm i\gamma_k, \quad k = 1, \dots, 100,$$

An Example

$$\zeta_2(s) = (1 - 2^{-s})\zeta(s)$$

Let us take 100 (pairs of conjugate) zeros of the zeta function,

$$\frac{1}{2} \pm i\gamma_k, \quad k = 1, \dots, 100,$$

and 201 zeros of $1 - 2^{-s}$,

$$0, \pm \frac{2\pi i}{\ln(2)}, \dots, \pm 100 \frac{2\pi i}{\ln(2)}$$

An Example

$$\zeta_2(s) = (1 - 2^{-s})\zeta(s)$$

Let us take 100 (pairs of conjugate) zeros of the zeta function,

$$\frac{1}{2} \pm i\gamma_k, \quad k = 1, \dots, 100,$$

and 201 zeros of $1 - 2^{-s}$,


$$0, \pm \frac{2\pi i}{\ln(2)}, \dots, \pm 100 \frac{2\pi i}{\ln(2)}$$

calculate corresponding 402 determinants of size 401 and normalize them getting numbers

$$\delta_{2,200,402,1}, \dots, \delta_{2,200,402,402}$$


$$\delta_{2,200,402,n} \text{ for } \zeta_2(s) = (1 - 2^{-s})\zeta(s)$$

$\delta_{2,200,402,n}$ for $\zeta_2(s) = (1 - 2^{-s})\zeta(s)$



Blue for $n \equiv 1 \pmod{2}$, Red for $n \equiv 0 \pmod{4}$, Green for $n \equiv 2 \pmod{4}$


$\delta_{2,200,402,n}$ for $\zeta_2(s) = (1 - 2^{-s})\zeta(s)$



Blue for $n \equiv 1 \pmod{2}$, Red for $n \equiv 0 \pmod{4}$, Green for $n \equiv 2 \pmod{4}$

$n < 120 \Rightarrow$ the δ 's are close to the coefficients of $(1 - 2^{-s})(1 - 2 \cdot 2^{-s})\zeta(s)$

$\delta_{2,200,402,n}$ for $\zeta_2(s) = (1 - 2^{-s})\zeta(s)$




Blue for $n \equiv 1 \pmod{2}$, Red for $n \equiv 0 \pmod{4}$, Green for $n \equiv 2 \pmod{4}$

$n < 120 \Rightarrow$ the δ 's are close to the coefficients of $(1 - 2^{-s})(1 - 2 \cdot 2^{-s})\zeta(s)$

$190 < n < 290 \Rightarrow$ the δ 's are close to the coefficients of $-2^{-s}(1 - 2 \cdot 2^{-s})\zeta(s)$

$\delta_{2,200,402,n}$ for $\zeta_2(s) = (1 - 2^{-s})\zeta(s)$



Blue for $n \equiv 1 \pmod{2}$, Red for $n \equiv 0 \pmod{4}$, Green for $n \equiv 2 \pmod{4}$


$n < 120 \Rightarrow$ the δ 's are close to the coefficients of $(1 - 2^{-s})(1 - 2 \cdot 2^{-s})\zeta(s)$

$190 < n < 290 \Rightarrow$ the δ 's are close to the coefficients of $-2^{-s}(1 - 2 \cdot 2^{-s})\zeta(s)$

$330 < n \Rightarrow$ the δ 's are very small


$$\log_{10} |\delta_{2,200,402,n} - 1| \text{ for } \zeta_2(s) = (1 - 2^{-s})\zeta(s)$$

$\log_{10} |\delta_{2,200,402,n} - 1|$ for $\zeta_2(s) = (1 - 2^{-s})\zeta(s)$



Blue for $n \equiv 1 \pmod{2}$, Red for $n \equiv 0 \pmod{4}$, Green for $n \equiv 2 \pmod{4}$

$\log_{10} |\delta_{2,200,402,n} - 1|$ for $\zeta_2(s) = (1 - 2^{-s})\zeta(s)$



Blue for $n \equiv 1 \pmod{2}$, Red for $n \equiv 0 \pmod{4}$, Green for $n \equiv 2 \pmod{4}$

Theorem of von Mangoldt

Теорема (Hans Carl Fridrich von Mangoldt [1895]).

$$\psi(x) = x - \sum_{\xi(\rho)=0} \frac{x^\rho}{\rho} - \sum_n \frac{x^{-2n}}{-2n} - \ln(2\pi)$$

Theorem of von Mangoldt

Теорема (Hans Carl Fridrich von Mangoldt [1895]).

$$\psi(x) = x - \sum_{\xi(\rho)=0} \frac{x^\rho}{\rho} - \sum_n \frac{x^{-2n}}{-2n} - \ln(2\pi)$$

$$\zeta(s) = \prod_{p - \text{простое}} \frac{1}{1 - p^{-s}}$$

Theorem of von Mangoldt

Теорема (Hans Carl Fridrich von Mangoldt [1895]).

$$\psi(x) = x - \sum_{\xi(\rho)=0} \frac{x^\rho}{\rho} - \sum_n \frac{x^{-2n}}{-2n} - \ln(2\pi)$$

$$\zeta(s) = \prod_{p - \text{простое}} \frac{1}{1 - p^{-s}}$$

$$\begin{aligned}\xi(s) &= \pi^{-\frac{s}{2}} (s-1) \Gamma\left(\frac{s}{2} + 1\right) \zeta(s) \\ &= \xi(0) \prod_{\xi(\rho)=0} \left(1 - \frac{s}{\rho}\right)\end{aligned}$$

Davenport–Heilbronn function

$$f(s) = \sum_{n=1}^{\infty} d(n)n^{-s}$$


where

$$d(n) = \begin{cases} 0, & \text{if } n \equiv 0 \pmod{5} \\ 1, & \text{if } n \equiv 1 \pmod{5} \\ \tau, & \text{if } n \equiv 2 \pmod{5} \\ -\tau, & \text{if } n \equiv 3 \pmod{5} \\ -1, & \text{if } n \equiv 4 \pmod{5} \end{cases}$$


and

$$\tau = \frac{-2 + \sqrt{10 - 2\sqrt{5}}}{-1 + \sqrt{5}}$$


Sieve of Eratosthenes for $f(s)$, red whenever $5|n$




Sieve of Eratosthenes for $f(s)$



Sieve of Eratosthenes for $f(s)$



Sieve of Eratosthenes for $f(s)$



Approximation of the zeta function

$$\Delta_N(s) = \sum_{n=1}^N \delta_{N,n} n^{-s}$$

Approximation of the zeta function

$$\begin{aligned}\Delta_N(s) &= \sum_{n=1}^N \delta_{N,n} n^{-s} \\ &\rightleftharpoons \sum_{n=1}^{\infty} (-1)^{n+1} n^{-s}\end{aligned}$$

Approximation of the zeta function

$$\begin{aligned}\Delta_N(s) &= \sum_{n=1}^N \delta_{N,n} n^{-s} \\ &\rightleftharpoons \sum_{n=1}^{\infty} (-1)^{n+1} n^{-s} \\ &= (1 - 2 \cdot 2^{-s}) \zeta(s)\end{aligned}$$

Approximation of the zeta function

$$\begin{aligned}\Delta_N(s) &= \sum_{n=1}^N \delta_{N,n} n^{-s} \\ &\stackrel{?}{=} \sum_{n=1}^{\infty} (-1)^{n+1} n^{-s} \\ &= (1 - 2 \cdot 2^{-s}) \zeta(s)\end{aligned}$$

$$\zeta(s) \stackrel{?}{\approx} \frac{\Delta_N(s)}{1 - 2 \cdot 2^{-s}}$$

Approximation of the zeta function

$$\begin{aligned}\Delta_N(s) &= \sum_{n=1}^N \delta_{N,n} n^{-s} \\ &\stackrel{?}{=} \sum_{n=1}^{\infty} (-1)^{n+1} n^{-s} \\ &= (1 - 2 \cdot 2^{-s}) \zeta(s)\end{aligned}$$

$$\zeta(s) \stackrel{?}{\approx} \frac{\Delta_N(s)}{1 - 2 \cdot 2^{-s}}$$

Approximation of the zeta function

$$\begin{aligned}
 \Delta_N(s) &= \sum_{n=1}^N \delta_{N,n} n^{-s} \\
 &\Leftarrow \sum_{n=1}^{\infty} (-1)^{n+1} n^{-s} \\
 &= (1 - 2 \cdot 2^{-s}) \zeta(s)
 \end{aligned}$$

$$\zeta(s) \stackrel{?}{\approx} \frac{\Delta_N(s)}{1 - 2 \cdot 2^{-s}}$$

s	$\left \frac{\Delta_{3001}(s)}{1 - 2 \cdot 2^{-s}} - \zeta(s) \right $
25	$4.2671 \dots \cdot 10^{-135}$
2	$3.9256 \dots \cdot 10^{-128}$
$1000i$	$4.4184 \dots \cdot 10^{-128}$
$\frac{1}{2} + 10i$	$1.0953 \dots \cdot 10^{-127}$
$-1 + 100i$	$3.6324 \dots \cdot 10^{-127}$
-25	$1.6415 \dots \cdot 10^{-126}$
$2 + 1000i$	$2.3063 \dots \cdot 10^{-125}$
$\frac{1}{2} + 1000i$	$3.9630 \dots \cdot 10^{-124}$
$-1 + 1000i$	$1.4867 \dots \cdot 10^{-118}$
$-10 + 1000i$	$8.2377 \dots \cdot 10^{-103}$
$\frac{1}{2} + 5000i$	$6.5116 \dots \cdot 10^{-64}$

Function $\nu_{N,L}(s)$

$$\Delta_N(s) = \sum_{n=1}^N \delta_{N,n} n^{-s} \leftrightharpoons \sum_{n=1}^{\infty} (-1)^{n+1} n^{-s} = (1 - 2 \cdot 2^{-s}) \zeta(s)$$

Function $\nu_{N,L}(s)$

$$\Delta_N(s) = \sum_{n=1}^N \delta_{N,n} n^{-s} \leftrightharpoons \sum_{n=1}^{\infty} (-1)^{n+1} n^{-s} = (1 - 2 \cdot 2^{-s}) \zeta(s)$$

$$\nu_N(s) = \frac{\Delta_N(s)}{\zeta(s)}$$

Function $\nu_{N,L}(s)$

$$\Delta_N(s) = \sum_{n=1}^N \delta_{N,n} n^{-s} \leftrightharpoons \sum_{n=1}^{\infty} (-1)^{n+1} n^{-s} = (1 - 2 \cdot 2^{-s}) \zeta(s)$$

$$\nu_N(s) = \frac{\Delta_N(s)}{\zeta(s)} = \frac{\sum_{n=1}^N \delta_{N,n} n^{-s}}{\sum_{n=1}^{\infty} n^{-s}}$$

Function $\nu_{N,L}(s)$

$$\Delta_N(s) = \sum_{n=1}^N \delta_{N,n} n^{-s} \leftrightharpoons \sum_{n=1}^{\infty} (-1)^{n+1} n^{-s} = (1 - 2 \cdot 2^{-s}) \zeta(s)$$

$$\nu_N(s) = \frac{\Delta_N(s)}{\zeta(s)} = \frac{\sum_{n=1}^N \delta_{N,n} n^{-s}}{\sum_{n=1}^{\infty} n^{-s}} = \sum_{n=1}^{\infty} \mu_{N,n} n^{-s}$$

Function $\nu_{N,L}(s)$

$$\Delta_N(s) = \sum_{n=1}^N \delta_{N,n} n^{-s} \leftrightharpoons \sum_{n=1}^{\infty} (-1)^{n+1} n^{-s} = (1 - 2 \cdot 2^{-s}) \zeta(s)$$

$$\nu_N(s) = \frac{\Delta_N(s)}{\zeta(s)} = \frac{\sum_{n=1}^N \delta_{N,n} n^{-s}}{\sum_{n=1}^{\infty} n^{-s}} = \sum_{n=1}^{\infty} \mu_{N,n} n^{-s} \leftrightharpoons 1 - 2 \cdot 2^{-s}$$

Function $\nu_{N,L}(s)$

$$\Delta_N(s) = \sum_{n=1}^N \delta_{N,n} n^{-s} \leftrightharpoons \sum_{n=1}^{\infty} (-1)^{n+1} n^{-s} = (1 - 2 \cdot 2^{-s}) \zeta(s)$$

$$\nu_N(s) = \frac{\Delta_N(s)}{\zeta(s)} = \frac{\sum_{n=1}^N \delta_{N,n} n^{-s}}{\sum_{n=1}^{\infty} n^{-s}} = \sum_{n=1}^{\infty} \mu_{N,n} n^{-s} \leftrightharpoons 1 - 2 \cdot 2^{-s}$$

$$\mu_{N,n} = \sum_{m|n} \mu\left(\frac{n}{m}\right) \delta_{N,n}$$

Function $\nu_{N,L}(s)$

$$\Delta_N(s) = \sum_{n=1}^N \delta_{N,n} n^{-s} \leftrightharpoons \sum_{n=1}^{\infty} (-1)^{n+1} n^{-s} = (1 - 2 \cdot 2^{-s}) \zeta(s)$$

$$\nu_N(s) = \frac{\Delta_N(s)}{\zeta(s)} = \frac{\sum_{n=1}^N \delta_{N,n} n^{-s}}{\sum_{n=1}^{\infty} n^{-s}} = \sum_{n=1}^{\infty} \mu_{N,n} n^{-s} \leftrightharpoons 1 - 2 \cdot 2^{-s}$$

$$\mu_{N,n} = \sum_{m|n} \mu\left(\frac{n}{m}\right) \delta_{N,n} \quad \delta_{N,n} = 0, \text{ if } n > N$$

Function $\nu_{N,L}(s)$

$$\Delta_N(s) = \sum_{n=1}^N \delta_{N,n} n^{-s} \leftrightharpoons \sum_{n=1}^{\infty} (-1)^{n+1} n^{-s} = (1 - 2 \cdot 2^{-s}) \zeta(s)$$

$$\nu_N(s) = \frac{\Delta_N(s)}{\zeta(s)} = \frac{\sum_{n=1}^N \delta_{N,n} n^{-s}}{\sum_{n=1}^{\infty} n^{-s}} = \sum_{n=1}^{\infty} \mu_{N,n} n^{-s} \leftrightharpoons 1 - 2 \cdot 2^{-s}$$

$$\mu_{N,n} = \sum_{m|n} \mu\left(\frac{n}{m}\right) \delta_{N,n} \quad \delta_{N,n} = 0, \text{ if } n > N$$

$$\zeta(s) = \frac{\Delta_N(s)}{\nu_N(s)}$$

Function $\nu_{N,L}(s)$

$$\Delta_N(s) = \sum_{n=1}^N \delta_{N,n} n^{-s} \leftrightharpoons \sum_{n=1}^{\infty} (-1)^{n+1} n^{-s} = (1 - 2 \cdot 2^{-s}) \zeta(s)$$

$$\nu_N(s) = \frac{\Delta_N(s)}{\zeta(s)} = \frac{\sum_{n=1}^N \delta_{N,n} n^{-s}}{\sum_{n=1}^{\infty} n^{-s}} = \sum_{n=1}^{\infty} \mu_{N,n} n^{-s} \leftrightharpoons 1 - 2 \cdot 2^{-s}$$

$$\mu_{N,n} = \sum_{m|n} \mu\left(\frac{n}{m}\right) \delta_{N,n} \quad \delta_{N,n} = 0, \text{ if } n > N$$

$$\zeta(s) = \frac{\Delta_N(s)}{\nu_N(s)} \quad \nu_{N,L}(s) = \sum_{n=1}^L \mu_{N,n} n^{-s}$$

Function $\nu_{N,L}(s)$

$$\Delta_N(s) = \sum_{n=1}^N \delta_{N,n} n^{-s} \leftrightharpoons \sum_{n=1}^{\infty} (-1)^{n+1} n^{-s} = (1 - 2 \cdot 2^{-s}) \zeta(s)$$

$$\nu_N(s) = \frac{\Delta_N(s)}{\zeta(s)} = \frac{\sum_{n=1}^N \delta_{N,n} n^{-s}}{\sum_{n=1}^{\infty} n^{-s}} = \sum_{n=1}^{\infty} \mu_{N,n} n^{-s} \leftrightharpoons 1 - 2 \cdot 2^{-s}$$

$$\mu_{N,n} = \sum_{m|n} \mu\left(\frac{n}{m}\right) \delta_{N,n} \quad \delta_{N,n} = 0, \text{ if } n > N$$

$$\zeta(s) = \frac{\Delta_N(s)}{\nu_N(s)} \quad \nu_{N,L}(s) = \sum_{n=1}^L \mu_{N,n} n^{-s}$$

$$\zeta(s) \stackrel{?}{\approx} \frac{\Delta_N(s)}{\nu_{N,L}(s)}$$

Function $\nu_{N,L}(s)$

$$\Delta_N(s) = \sum_{n=1}^N \delta_{N,n} n^{-s} \leftrightharpoons \sum_{n=1}^{\infty} (-1)^{n+1} n^{-s} = (1 - 2 \cdot 2^{-s}) \zeta(s)$$

$$\nu_N(s) = \frac{\Delta_N(s)}{\zeta(s)} = \frac{\sum_{n=1}^N \delta_{N,n} n^{-s}}{\sum_{n=1}^{\infty} n^{-s}} = \sum_{n=1}^{\infty} \mu_{N,n} n^{-s} \leftrightharpoons 1 - 2 \cdot 2^{-s}$$

$$\mu_{N,n} = \sum_{m|n} \mu\left(\frac{n}{m}\right) \delta_{N,n} \quad \delta_{N,n} = 0, \text{ if } n > N$$



$$\zeta(s) = \frac{\Delta_N(s)}{\nu_N(s)} \quad \nu_{N,L}(s) = \sum_{n=1}^L \mu_{N,n} n^{-s}$$

$$\zeta(s) \stackrel{?}{\approx} \frac{\Delta_N(s)}{\nu_{N,L}(s)} = \frac{\sum_{n=1}^N \delta_{N,n} n^{-s}}{\sum_{n=1}^L \mu_{N,n} n^{-s}}$$

Calculation of $\zeta(s)$ at $s = \frac{1}{4} + 1000i$ for $N = 3001$

L	$\mu_{N,L}$	$\left \zeta(s) - \frac{\Delta(s)}{\nu_{N,L}(s)} \right $
2	$-2 + 1.43 \dots \cdot 10^{-127}$	$2.24128 \dots \cdot 10^{-127}$
3	$-2.14787 \dots \cdot 10^{-127}$	$1.57968 \dots \cdot 10^{-299}$
4	$-1.62673 \dots \cdot 10^{-299}$	$4.85859 \dots \cdot 10^{-448}$
5	$+5.29034 \dots \cdot 10^{-448}$	$1.00748 \dots \cdot 10^{-569}$
6	$-1.14817 \dots \cdot 10^{-569}$	$1.83153 \dots \cdot 10^{-672}$
7	$+2.16930 \dots \cdot 10^{-672}$	$3.15150 \dots \cdot 10^{-756}$
8	$-3.85941 \dots \cdot 10^{-756}$	$2.34266 \dots \cdot 10^{-829}$
9	$-2.95462 \dots \cdot 10^{-829}$	$3.17791 \dots \cdot 10^{-891}$
10	$+4.11503 \dots \cdot 10^{-891}$	$6.45307 \dots \cdot 10^{-946}$
11	$-8.55748 \dots \cdot 10^{-946}$	$6.55682 \dots \cdot 10^{-994}$
12	$+8.88627 \dots \cdot 10^{-994}$	$1.00011 \dots \cdot 10^{-1036}$
13	$+1.38282 \dots \cdot 10^{-1036}$	$2.32048 \dots \cdot 10^{-1074}$
14	$-3.26844 \dots \cdot 10^{-1074}$	$1.18994 \dots \cdot 10^{-1108}$

Numbers $\log_{10} |\mu_{3001,n}|$



$$\mu_{N,n} = \sum_{m|n} \mu\left(\frac{n}{m}\right) \delta_{N,n}$$